

General static black holes in matter

K.A. Bronnikov¹

*Center for Gravitation and Fundamental Metrology, VNIIMS, 46 Ozyornaya St., Moscow 119361, Russia;
Institute of Gravitation and Cosmology, PFUR, 6 Miklukho-Maklaya St., Moscow 117198, Russia*

Oleg B. Zaslavskii²

Astronomical Institute of Kharkov V.N. Karazin National University, 35 Sumska St., Kharkov, 61022, Ukraine

For arbitrary static space-times, it is shown that an equilibrium between a Killing horizon and matter is only possible for some discrete values of the parameter $w = p_1/\rho$, where ρ is the density and p_1 is pressure in the direction normal to the horizon. In the generic situation of a simple (non-extremal) horizon and the slowest possible density decrease near the horizon, this corresponds to $w = -1/3$, the value known for a gas of disordered cosmic strings. An admixture of “vacuum matter”, characterized by $w = -1$ and nonzero density at the horizon, is also admitted. This extends the results obtained previously for static, spherically symmetric space-times. A new feature as compared to spherical symmetry is that higher-order horizons can exist in the absence of vacuum matter if the horizon is a surface of zero curvature, which can occur, e.g., in cylindrically symmetric space-times.

PACS: 04.70.Bw, 04.20.Cv, 04.40.Nr

1 Introduction

It is generally believed that, in the course of gravitational collapse, as a black hole is being formed, nonspherical perturbations die out. However, under real astrophysical conditions, black holes are surrounded by different kinds of celestial bodies and distributed matter (stars in double and multiple stellar systems, gas clouds etc.) Even if a black hole can be regarded static in a good approximation, these bodies distort its spherical shape, and not always such deflections are small. On the other hand, distributed matter can either be in equilibrium with the black hole or fall on it. Conditions of such equilibrium are of substantial interest.

It is also worth noting that the generic properties of Killing horizons (not necessarily spherically symmetric) play an important role in the (not yet well understood) relationship between the horizon symmetry and black hole entropy (see, e.g., [1] and references therein). A discussion of black hole entropy in connection with statistical properties of the surrounding matter also requires a knowledge of which kinds of matter can be in equilibrium with the horizon.

In our previous paper [2], we studied the equilibrium conditions between macroscopic matter and static, spherically symmetric black holes. Using near-horizon expansions in the Einstein equations

¹e-mail: kb20@yandex.ru

²e-mail: ozaslav@kharkov.ua

and the conservation law for matter, we found that only some discrete values of the parameter $w = p_r/\rho$ (the radial pressure to density ratio) are compatible with such equilibrium, and its generic values are $w = -1$ (which is well known [1, 3]) and $w = -1/3$ (which corresponds to a gas of cosmic strings if this matter is isotropic); normal ($w \geq 0$) or phantom ($w < -1$) matter is excluded.

In the present paper we extend this approach to general static black holes. It turns out that almost all conclusions, made previously for the case of spherical symmetry, remain valid when this requirement is relaxed.

2 Basic equations

We consider the generic static metric which can be written, in terms of the so-called Gaussian coordinates, in the form³

$$ds^2 = N^2 dt^2 - dl^2 - \gamma_{ab} dx^a dx^b, \quad a, b = 2, 3, \quad (1)$$

where N and elements of the 2-metric γ_{ab} are functions of l, x^2, x^3 . Horizons, if any, are located at surfaces where $N = 0$.

Let us assume that the stress-energy tensor (SET) of matter has the components $T_0^0 = \rho$, $T_1^1 = p_1$ ($x^1 = l$), $T_0^i = 0$ ($i = 1, 2, 3$). Then the l component of the conservation law $\nabla_\nu T_\mu^\nu = 0$ gives

$$p_1' + (p_1 + \rho) \frac{N'}{N} + \left(p_1 \frac{\gamma'}{2\gamma} + K_{ab} T^{ab} \right) = 0, \quad (2)$$

where the prime denotes $\partial/\partial l$, $\gamma = \det(\gamma_{ab})$, and

$$K_{ab} = -\frac{1}{2} \frac{\partial \gamma_{ab}}{\partial l} \quad (3)$$

is the extrinsic curvature tensor of the surfaces $t = \text{const}$, $l = \text{const}$ showing how they are embedded in the outer three-space.

Using the explicit expressions for the Einstein tensor given in [1], we can write two components of the Einstein equations as follows:

$$G_1^1 - G_0^0 = \frac{\Delta_2 N}{N} + K' - \frac{N'}{N} K - K_{ab} K^{ab} = 8\pi(p_1 + \rho), \quad (4)$$

$$G_1^1 = -\frac{1}{2} R_{||} - \frac{1}{2} K_{ab} K^{ab} + \frac{1}{2} K^2 + \frac{\Delta_2 N}{N} - \frac{K N'}{N} = 8\pi p_1, \quad (5)$$

where $K = \text{tr } K = \gamma^{ab} K_{ab}$, $R_{||}$ is the scalar curvature of the two-surface $t = \text{const}$, $l = \text{const}$, and Δ_2 is the two-dimensional Laplacian corresponding to the metric γ_{ab} on such a surface.

³We use the units $c = \hbar = G = 1$.

3 Horizons and their neighborhood

Considering a two-dimensional (t, l) section of our manifold at fixed values of the coordinates x^a , we can introduce the so-called quasiglobal coordinate u by the relation $du = N(l)dl$. (Note that if we try to make such a transformation in four dimensions with $N = N(l, x^a)$, we shall, in general, obtain nonzero mixed terms $g_{1a}du dx^a$ in the linear element. Therefore, to obtain correct relations, we fix x^a assuming that all derivatives in x^a in our equations have already been calculated.) We thus have at fixed x^a

$$ds_2^2 = A(u)dt^2 - du^2/A(u), \quad (6)$$

where $A(u) \equiv N^2(l)$. This coordinate has the following important properties (see, e.g. [4–6]): it always takes a finite value $u = u_h$ at a Killing horizon where $A(u) = 0$;⁴ moreover, near a horizon, the increment $u - u_h$ is a multiple (with a nonzero constant factor) of the corresponding increments of manifestly well-behaved Kruskal-type null coordinates, used for analytic extension of the metric across the horizon. Thus a horizon corresponds to a regular zero of $A(u)$, i.e.,

$$A(u) \equiv N^2(l) \sim (u - u_h)^n, \quad (7)$$

where $n \in \mathbb{N}$ is the order of the horizon. A simple, Schwarzschild-like horizon corresponds to $n = 1$, an extremal one (as in extremal Reissner-Nordström black holes) to $n = 2$, and horizons with $n > 2$ are called ultraextremal.

Thus all metric functions should be analytic (or smooth at least up to derivatives of a certain order $s \geq 2$) functions of u at $u = u_h$. In particular, this applies to γ_{ab} which behave as scalars in the (t, u) subspace. Then

$$K_{ab} = -\frac{1}{2} \frac{\partial \gamma_{ab}}{\partial l} = -\frac{1}{2} \sqrt{A} \frac{d\gamma_{ab}}{du} \sim (u - u_h)^{n/2}; \quad (8)$$

here and henceforth expressions in terms of u are valid at fixed x^a . Thus $K_{ab} = O(N) \rightarrow 0$ at the horizon.

Furthermore, assuming a Taylor expansion for $A(u)$ near $u = u_h$, it is easy to find the near-horizon behavior of $N(l)$: in accord with [1, 7, 8], we obtain

$$n = 1 : \quad l \rightarrow 0, \quad N(l) = \varkappa_1 l + \varkappa_3 l^3 + \dots; \quad (9)$$

$$n = 2 : \quad l \rightarrow \infty, \quad N(l) = B_1 e^{-bl} + B_2 e^{-2bl} + \dots; \quad (10)$$

$$n > 2 : \quad l \rightarrow \infty, \quad N(l) = C_n e^{-n/(n-2)} + C_{n+2} e^{-(n+2)/(n-2)} + \dots, \quad (11)$$

where the quantities $\varkappa_i$, b , B_i , C_i are, in general, functions of x^2 , x^3 , since these expansions have been obtained for fixed x^a .

It has been shown, however [7, 8], that the quantities $\varkappa_1$, B_1 and C_n are constants. For simple horizons, $\varkappa_1 = \varkappa$ is nothing else but the surface gravity, and its constancy along the horizon is the

⁴In principle, u can take an infinite value at a candidate horizon where $A \rightarrow 0$, but then, as one can check, the canonical parameter of the geodesics also tends to infinity, so that the space-time is already geodesically complete and no continuation is required. Such cases (“remote horizons”) are possible in principle, but we will not discuss them here.

content of the so-called zeroth law of black hole thermodynamics [3]. For higher-order horizons ($n \geq 2$), the constancy of B_1 and C_n has been proved in [7, 8] from the horizon regularity requirement: if one admits their x^a dependence, one obtains the so-called truly naked horizons, at which some curvature components in the freely falling reference frame are infinite even though all algebraic curvature invariants remain finite.

Due to x^a -independence of the first terms in the expansions (9)–(11), it is possible to introduce the quasiglobal coordinate u in our four-dimensional space-time as a whole, not only on its (t, l) -sections, at least near the horizons, and preserve the whole reasoning connected with regularity in terms of u . Thus, e.g., for simple horizons we have

$$l = \sqrt{\varkappa/2} \sqrt{u - u_h} [1 + l_3(x^a)(u - u_h) + \dots],$$

so that an x^a -dependence is absent in the main approximation with respect to $u - u_h$; in the same approximation, we can also neglect the possible x^a -dependence of u_h (though, it does not affect our reasoning in any way).

4 Matter compatible with a horizon

From the expressions (9)–(11) it follows that N'/N is zero at the horizon for $n = 1$ and $n > 2$, and it has there a finite value for $n = 2$. From the constancy of $\varkappa$, B_1 and C_1 it also follows that $\Delta_2 N/N \rightarrow 0$ as $N \rightarrow 0$. Together with (8) this implies $G_1^1 - G_0^0 = 0$ at the horizon, whence it follows

$$p_1 + \rho = 0 \tag{12}$$

at the horizon, in accord with [1] (though our proof is different from that in [1]).

Let us assume that (at least near the horizon) the equation of state of matter takes the form

$$p_1 \approx w\rho \tag{13}$$

with $w = \text{const}$. We also suppose that

$$|T_a^b|/\rho < \infty. \tag{14}$$

In the conservation law (2), the ratio of the third term to the second one is of the order $(\max K_{ab})N/N'$. For simple horizons ($n = 1$) $N' > 0$, and in the extremal case ($n = 2$) $N' \sim N$ whereas $K_{ab} \sim N$ (at most). Thus in both these cases we can safely neglect the third term in (2). In the ultraextremal case ($n > 2$) $N/N' \sim l \rightarrow \infty$; however, due to (11) the ratio in question behaves as $l^{-2/(n-2)} \rightarrow 0$ as $l \rightarrow \infty$, so that in this case the third term in (2) is also negligible as compared with the second one.

With the third term neglected, Eqs. (2) and (13) lead to

$$\rho \sim N^{-(1+w)/w}. \tag{15}$$

Thus the admissible range of w , in which $\rho < \infty$ at the horizon, is $-1 \leq w < 0$, where $w = -1$ corresponds to the so-called vacuum fluid [9] whose definitive property is that in a certain spatial

direction x^1 there holds the relation (12). It generalizes the notion of a spherically symmetric vacuum fluid (see also [2] and references therein) and certainly also includes the cosmological constant as a special case.

For other values of w , we have $\rho \rightarrow 0$ as $N \rightarrow 0$.

On the other hand, as before, we can consider the Einstein equations (4) and (5) at fixed values of the coordinates x^a , and then, since all metric functions should depend analytically (or sufficiently smoothly) on the quasiglobal coordinate u , the same is true for the right-hand side of these equations, which, for $w > -1$, means that $\rho \sim (u - u_h)^k$ where k is a positive integer. Comparing this condition with (7) and (15), we obtain

$$w = -\frac{n}{n + 2k}, \quad (16)$$

thus reproducing the result of [2], obtained there for spherically symmetric configurations. Thus matter which can be in equilibrium with static horizons of any shape should have an equation of state with the parameter w from the discrete set of values (16), where n is the order of the horizon and k characterizes the near-horizon behavior of the density.

We can also admit a non-interacting mixture of our matter and a vacuum fluid with the property $\rho_{(\text{vac})} + p_{1(\text{vac})} = 0$. The conservation law (2) then holds for each of them separately, and nothing is changed in the above reasoning.

One more restriction follows from Eq. (5). Indeed, at the horizon itself it gives

$$\frac{1}{2}R_{\parallel} + K_1 = 8\pi\rho_{(\text{vac})}, \quad K_1 := \lim_{\text{hor}} \frac{KN'}{N}. \quad (17)$$

Generically (if K does not vanish quicker than N when approaching the horizon), we have $K_1 \sim (u - u_h)^{n-1}$.

Further conclusions depend on the presence of a vacuum fluid.

1. $\rho_{(\text{vac})} = 0$. In the general case that the horizon is curved, $R_{\parallel} \neq 0$, according to (17), we must have $K_1 \neq 0$, so that the horizon must be simple, $n = 1$. Higher-order horizons ($n \geq 2$) are only possible if $R_{\parallel} = 0$. In particular, they cannot exist in spherically symmetric space-times, in accord with [2], but they can appear in cylindrically symmetric black holes since the corresponding two-dimensional cross-section is flat for them.

2. $\rho_{\text{vac}} \neq 0$. Eq. (17) admits any order of the horizon but, if it is higher-order, $K_1 = 0$, hence flat horizons ($R_{\parallel} = 0$) are now impossible. For $n \geq 2$, Eq. (17) gives a direct relationship between the horizon curvature and the vacuum fluid density

$$R_{\parallel} = 16\pi\rho_{(\text{vac})}.$$

Thus, without knowing any exact solutions, we have obtained certain restrictions on the possible horizon geometry depending on the existence and properties of the vacuum fluid.

5 Concluding remarks

We have shown that the main result of [2], obtained there for Killing horizons in static, spherically symmetric space-times, also holds for horizons in general static space-times: an equilibrium between

a horizon and matter is only possible for some particular values of the parameter $w = p_1/\rho$ given by Eq. (16). In the generic situation of a simple (non-extremal) horizon and the slowest possible density decrease near the horizon, this corresponds to $w = -1/3$, the value characteristic of a gas of disordered cosmic strings.

A new feature as compared to spherical symmetry is that, in the absence of vacuum matter, higher-order horizons are possible provided the horizon as a 2-surface has zero scalar curvature $R_{\parallel}$. In particular, this is true for cylindrical black holes. If $\rho_{(\text{vac})} \neq 0$, the sign of $\rho_{(\text{vac})}$ should coincide with that of $R_{\parallel}$. In particular, $\rho_{\text{vac}} < 0$, forbidden in the spherical case, is compatible with a horizon with hyperbolic geometry. Thus matter with certain negative values of w can contain horizons with non-spherical topologies (cf. [10]).

In our reasoning, which has been entirely local and relied on near-horizon expansions, we did not assume any particular equation of state for matter and even did not restrict the behaviour of the transverse pressure except for its regularity requirement. Moreover, no assumptions on the asymptotic properties of the space-time manifold were made. In this sense, our conclusions are model-independent.

References

- [1] A. J. M. Medved, D. Martin and M. Visser, *Class. Quantum Grav.* **21**, 3111 (2004).
- [2] K. A. Bronnikov and O. B. Zaslavskii, *Phys. Rev. D* **78**, 021501(R) (2008).
- [3] V. P. Frolov and I. D. Novikov, *Black Hole Physics: Basic Concepts and New Developments* (Kluwer Academic, Boston, 1998).
- [4] K. A. Bronnikov, *Phys. Rev. D* **64**, 064013 (2001).
- [5] K. A. Bronnikov, G. Clément, C. P. Constantinidis, and J. C. Fabris, *Grav. Cosmol.* **4**, 128 (1998); *Phys. Lett. A* **243**, 121 (1998).
- [6] K. A. Bronnikov, E. Elizalde, S.D. Odintsov and O. B. Zaslavskii, *Phys. Rev. D* **78**, 064049 (2008).
- [7] O. B. Zaslavskii, *Phys. Rev. D* **76**, 024015 (2007).
- [8] V. Pravda and O. B. Zaslavskii, *Class. Quantum Gravity* **22**, 5053 (2005).
- [9] I.G. Dymnikova, *Gen. Rel. Grav.* **24**, 235 (1992).
- [10] D.R. Brill, J. Louko and Peldan, *Phys. Rev. D* **56**, 3600 (1997);
L. Vanzo, *Phys. Rev. D* **56**, 6475 (1997);
J.P.S. Lemos, *Phys. Rev. D* **57**, 4600 (1998).